

***COMMENT ON “SEASONAL VARIATION OF THE PHYSICAL PROPERTIES
OF MARINE BOUNDARY LAYER CLOUDS OFF THE CALIFORNIA
COAST” BY LIN ET AL.***

Yangang Liu

Accepted for publication in
J. Climate

February 2010

Environmental Sciences Department/Atmospheric Sciences Division

Brookhaven National Laboratory

P.O. Box 5000

Upton, NY 11973-5000

www.bnl.gov

Notice: This manuscript has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-98CH10886 with the U.S. Department of Energy. The publisher by accepting the manuscript for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes.

This preprint is intended for publication in a journal or proceedings. Since changes may be made before publication, it may not be cited or reproduced without the author's permission.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

By combining data from multiple sources, Lin et al. (2009) have lately performed a rather comprehensive comparison of the seasonal differences in several macrophysical properties of marine boundary layer (MBL) clouds observed off the California coast (e.g., cloud fraction, liquid water path, cloud-top and cloud-base heights, cloud thickness, inversion strength, lifted condensation level, and the degree of decoupling). They found that most differences between the summer (JJA) and winter (DJF) seasons can be explained by the characteristics of lower tropospheric stability (LTS, Slingo 1980; Klein, 1997) and the deepening-warming-decoupling hypothesis proposed in Bretherton and Wyant (1997). Although this work certainly constitutes an excellent contribution to the understanding and parameterization of MBL clouds by considering multiple variables together, it is primarily confined to macrophysical properties, only with a brief mentioning of the cloud droplet effective radius. Below I will demonstrate that credible microphysical information can in fact be inferred from the same datasets provided in Lin et al., which further suggests at least equally strong summer-winter differences in microphysical properties and a plausible microphysical effect. Note that “microphysical” is here used in a general sense to denote any variables that can be derived from a local cloud droplet size distribution without involving cloud geometrical properties (e.g., liquid water content, droplet concentration, and effective radius).

The first microphysical quantity is the cloud-layer mean liquid water content (L). In their Fig. 5b and 5f, Lin et al. compared liquid water path (LWP) and cloud thickness (H) observed during the summer and winter seasons, and concluded that the summer-winter difference in LWP is much larger than that in the cloud thickness. Because LWP can be regarded as a product of H and L

$$\text{LWP} = \text{HL}, \quad (1)$$

L is readily obtained from LWP and H by use of the variant of (1): $L = \text{LWP}/H$. Accordingly, a large summer-winter difference in L is anticipated from the distinct contrast between the summer-winter differences in LWP and H. This point can be clearly seen in Fig 1a, which contrasts the variations of the summertime and wintertime L with the distance from the California coast. The panels 1b and 1c are reproduction of their Fig. 5b and 5e for the convenience of comparison. The higher summertime L is likely related to the larger dq_{sat}/dz associated with the corresponding lower cloud-base height as reported in Lin et al; q_{sat} is the saturation mixing ratio and adiabatic liquid water content is approximately proportional to dq_{sat}/dz at a given height z (Albrecht et al., 1990). In addition to the expected summer-winter difference, L also exhibits a unique variation with the distance from the coast: In wintertime, the farther away from the California coast, the higher the L value; but the opposite appears true in summertime. But, the physics behind the converging trend of the summertime and wintertime L from the coast to the ocean is elusive at present.

The second microphysical quantity is the cloud droplet number concentration (N). Without providing a detailed analysis, Lin et al mentioned that, “Retrievals from CERES-MODIS, however, have shown that the effective radius is smaller in JJA than in DIF”. Crucial information on N can be further inferred from the results of L and effective radius by examining the relationship between the three microphysical variables. Without loss of generality, effective radius (r_e) is related to L and N by the general power-law relationship (Liu and Hallett 1997; Liu and Daum 2000; Liu et al., 2008)

$$r_e = a_e \left(\frac{L}{N} \right)^{b_e}, \quad (2)$$

where a_e and b_e are two empirical coefficients that have been often assumed as constant, especially in remote sensing of cloud properties (e.g., Han et al., 1998; Boers et al., 2006).

With the data on L and r_e , N can be readily estimated using (2) with a given pair of a_e and b_e [see more discussion on the two coefficients in Boer et al. (2006) and Liu et al. (2008)]. In fact, a few studies have attempted to retrieve N from satellite measurements based on equations similar to (2) (Han et al., 1998; Szczodrak et al., 2001; Schuller et al., 2003; Boers et al., 2006). Even without knowing the specific values of N , some salient feature of the summer-winter difference in N can be deduced by mathematical analysis of (2). Differentiation of (2) yields the equation that describes the fractional differences in N , L , and effective radius

$$\frac{\Delta N}{N} = \frac{\Delta L}{L} - \frac{1}{b_e} \frac{\Delta r_e}{r_e} + \frac{1}{b_e} \frac{\Delta a_e}{a_e}, \quad (3a)$$

where Δ denotes the difference between the summer- and winter-clouds. To the first order approximation, it can be assumed that the two coefficients a_e and b_e are same during the summer and winter seasons. Under this assumption, (3a) is simplified as

$$\frac{\Delta N}{N} = \frac{\Delta L}{L} - \frac{1}{b_e} \frac{\Delta r_e}{r_e} \quad (3b)$$

Equation (3b) clearly reveals that when the summertime r_e is smaller than that of the winter cloud, or $\Delta r_e < 0$, the difference in L not only means a difference in N , but more interesting is that the summer-winter difference in N is larger than that in L , or

$$\frac{\Delta N}{N} > \frac{\Delta L}{L}. \quad (3c)$$

More specific information on the relationship between the relative differences can be obtained by examining two special values of b_e . The first case is the commonly used one of $b_e = 1/3$. For this case, the relative summer-winter difference in N is given by

$$\frac{\Delta N}{N} = \frac{\Delta L}{L} - 3 \frac{\Delta r_e}{r_e}. \quad (4a)$$

Despite its popularity, $b_e = 1/3$ holds only if the spectral shape of the cloud droplet size distribution remains unchanged when N and L vary. However, recent studies have shown that some physical processes simultaneously affect L , N , and the spectral shape (Liu and Daum, 2002; Yum and Hudson, 2005; Liu et al., 2006; Peng et al, 2007; Hsieh et al., 2009). By analyzing a collection of cloud data obtained during several field campaigns, Liu et al (2008) obtained $b_e = 0.19$, smaller than the commonly use $b_e = 1/3$ because of concurrent increases of N and relative dispersion of the cloud droplet size distribution and spectral narrowing of adiabatic condensational growth. For the case of $b_e = 0.19$, we have

$$\frac{\Delta N}{N} = \frac{\Delta L}{L} - 5.26 \frac{\Delta r_e}{r_e} \quad (4b)$$

Comparison of (4b) with (4a) shows that consideration of the spectral shape effect results in an even larger difference in N .

The third microphysical quantity is the drizzle rate, which can be examined using the diagnostic relationship between cloud-base drizzle rate P , LWP and N found from surface-based remote sensing (Wood et al. 2008),

$$P = 0.37 \left(\frac{LWP}{N} \right)^{1.75}, \quad (5)$$

where P is unit of mmd^{-1} , LWP in gm^{-2} , and N in cm^{-3} . A combination of (1), (2), and (5) leads to

$$P = 0.37 \left(\frac{LWP}{N} \right)^{1.75} = 0.37 a_e^{-1.75/b_e} r_e^{1.75/b_e} H^{1.75} \quad (6)$$

Evidently, P can be estimated from LWP and N using the first identity, or from r_e and H using the second identity. Using the second identity also needs to assume the values of a_e and b_e . Again, useful information can be inferred from what Lin et al. provided without knowing the concrete values of N and r_e , Lin et al. reported that compared to the winter cloud, the summertime cloud has a smaller r_e but approximately same H . Together with (6), this finding suggests that the summertime cloud has a smaller drizzle rate than the wintertime cloud.

It is interesting to note that the totality of the microphysical summer-winter differences inferred from what Lin et al. provided ---- the summertime cloud has a higher L and N but lower r_e and P ---- resembles the majority of the microphysical differences between closed and open convective cells in MBL clouds, and/or between the so-called pockets of open cells (POCs) and their surrounding solid clouds. In search for understanding cellular cloud structure, there have been increasing number of studies on the connection between the cellular structure and microphysics via drizzles (e.g., Stevens et al., 2005; Petters et al., 2006; Sharon et al., 2006; Wood et al. 2008). These studies have reported compelling observational evidence that the microphysical characteristics in solid decks or closed cells differ substantially from those of POCs or open cells, with open cells or POCs being associated with lower N , but larger r_e and P . Numerical simulations (Savic-Jovicic and Stevens 2008; Xue et al. 2008; Wang and Feingold 2009) have confirmed these observations, and further suggested that enhanced precipitation plays a critical role in the formation and evolution of open cells, and evaporation of raindrops generate a dynamic response that promotes, organizes and sustains open cell structures. Lower CCN/aerosol

concentrations have also been found to be associated with open cells and POCs, implying potential aerosol influences. Although it cannot be conclusive without supporting CCN/aerosol data, the remarkable microphysical similarity between the summer-winter microphysical differences inferred from Lin et al. and those between the closed (solid decks) and open cells (POCs) is certainly indicative of a microphysical mechanism for winter clouds that is in action for open cells and POCs. The summer-winter differences in the microphysical properties are also consistent with the dominant mechanisms proposed for aerosol indirect effects: an increase in aerosol loading leads to a higher N but a smaller effective radius (Twomey 1974), and less drizzles but higher L and LWP (Albrecht 1989).

Lin et al. pointed out that the seasonal variations of the macrophysical cloud properties from summer to winter resemble the downstream stratocumulus-to-cumulus transition of MBL clouds, and that the “deepening-warming-decoupling” mechanism proposed by Bretherton and Wyant (1997) can explain the summer-winter cloud differences when the warming of the sea surface temperature is relative to the temperature of the free troposphere. Together with the plausible microphysical mechanism, the question arises as to which mechanism (the macrophysical discussed in Lin et al or the microphysical mechanism discussed here) is more important in determining the seasonal differences. Of course, macrophysics and microphysics are likely the two sides of the same coin, and the seasonal differences may stem from interwoven actions of both macrophysical and microphysical mechanisms via multiscale interactions/feedbacks.

Responding to my original comment, Lin et al have performed the microphysical analyses suggested above, and their results largely support the expected microphysical differences outlined above (see their response for details). Deeper insights can be obtained

by examining their new results. Table 1 juxtaposes the main macrophysical and microphysical quantities for the summertime and wintertime clouds and compared their relative differences defined as (summertime – wintertime)/summertime. The six variables that have the largest relative summer-winter differences are: P (-255%), cloud-base height (-121%), N (72%), cloud-top height (-72%), L (42%), and r_e (-28%). Note that the difference in liquid water path (43%) is not listed because it primarily reflects the difference in liquid water content as the wintertime and summertime clouds have similar cloud thickness. These composite results seem to support joint roles of macrophysics and microphysics via drizzles, as a higher cloud-base in the winter cloud tends to associate with lower droplet concentration and liquid water content. An analytical steady-state formulation also confirms the essential role of N in determining P (Wood 2009, private communication).

The new results also indicate that like L, the changes of N, r_e , and P with the distances from the California coast all exhibit a converging trend (Fig. 1, Lin et al., 2009). Figure 2 further compares the relative summer-winter differences in the four microphysical properties as a function of the distance away from the California coast. As for the mean discussed above, Fig. 2 shows that P and N have the largest seasonal differences from the coast to the ocean, providing additional support for the crucial role of N in determining P. However, it has proven difficult to tell whether a smaller N (aerosols) causes a higher P first or is caused by a larger P because of the positive feedback loop between N and P. The converging behaviors of the microphysical differences shed further insight on this issue: The gradual decreases of all the microphysical differences with increasing distances from the California coast suggests the importance of coastal proximity and thus aerosols in shaping the summer-winter differences.

It has been long recognized that MBL clouds are highly coupled systems with complex interactions between thermodynamics, dynamics, radiation and microphysics, and growing efforts have been recently devoted to understanding the co-workings of macrophysics and microphysics (Kubar et al., 2009; Wood et al., 2009). Nevertheless, a full theoretical framework for such highly coupled systems is still posing a challenge. This is especially true for developing an adequate representation of MBL clouds in climate models where cloud macrophysics (e.g., cloud fraction) and microphysics (e.g., aerosol effects) are often treated separately. More comprehensive analyses and idealized numerical simulations should be essential for addressing these intriguing issues.

Acknowledgement: This work is supported by the U.S. Department of Energy's Atmospheric Science Research (ASR) Program and the Earth System Modeling (ESM) Program.

References

- Albrecht, B., 1989: Aerosols, cloud microphysics, and fractional cloudiness. *Science*, **245**, 1227-1230.
- Boers, R., J. R. Acarreta, and J. L. Gras, 2006: Satellite monitoring of the first indirect aerosol effect: retrieval of the droplet concentration of water clouds. *J. Geophys. Res.*, **111**, D22208, doi:10.1029/2005JD006838.
- Bretherton, C. S., and M. C. Wyant, 1997: Moisture transport, lower-tropospheric stability, and decoupling of cloud-topped boundary layers. *J. Atmos. Sci.*, **54**, 148-167.
- Han, Q., W. B. Rossow, J. Chou, and R. M. Welch, 1998: Global variation of column droplet concentration in a low-level clouds. *Geophys. Res. Lett.*, **25**, 1419-1422.
- Hsieh, W. C. and coauthors, 2009: Parameterization of cloud droplet size distributions: comparison with parcel models and observations. *J. Geophys. Res.*, **114**, D11205, doi:10.1029/2008JD011387.
- Kubar, T. L., D. L. Hartmann, and R. Wood, 2009: Understanding the importance of microphysics and macrophysics for warm rain in marine low clouds. Part I: Satellite observations. *J. Atmos. Sci.*, **66**, 2953-2972.
- Lin, W., M. Zhang, and N. G. Loeb, 2009: Seasonal variation of the physical properties of marine boundary layer clouds off the California coast. *J. Climate*, **22**, 2624-2638.
- Liu, Y. and P. H. Daum, 2000: Spectral dispersion of cloud droplet size distributions and the parameterization of cloud droplet effective radius. *Geophys. Res. Lett.* **27**, 1903-1906.
- Liu, Y., and P. H. Daum, 2002: Indirect warming effect from dispersion forcing. *Nature*, **419**, 580-581.

- Liu, Y., P. H. Daum, H. Guo, and Y. Peng, 2008: Dispersion bias, dispersion effect, and the aerosol-cloud conundrum. *Environ. Res. Lett.*, **3**, 04502, doi:10.1088/1748-9326/3/4/045021.
- Liu, Y., P. H. Daum and S. Yum, 2006: Analytical expression for the relative dispersion of the cloud droplet size distribution. *Geophys. Res. Lett.*, **33**, L02810, doi:10.1029/2005GL024052.
- Liu, Y., and J. Hallett, 1997: The "1/3" power-law between effective radius and liquid water content. *Quart. J. Roy. Meteor. Soc.* **123**, 1789-1795.
- Peng, Y., U. Lohmann, R. Leaith, and M. Kulmala, 2007: An investigation into the aerosol dispersion effect through the activation process in marine stratus clouds. *J. Geophys. Res.*, **112**, D1117, doi:10.1029/2006JD007401.
- Savic-Jovcic, V., and B. Stevens, 2008: The structure and mesoscale organization of precipitating stratocumulus. *J. Atmos. Sci.*, **65**, 1587-1605.
- Schuller, L., J. Brenguier, and H. Pawlowska, 2003: Retrieval of microphysical, geometrical, and radiative properties of marine stratocumulus from remote sensing. *J. Geophys. Res.*, **108**, D8631, doi:10.1029/2002JD002680.
- Szczodrak, M., P. H. Austin, and P. B. Krummel, 2001: Variability of optical depth and effective radius in marine stratocumulus clouds. *J. Atmos. Sci.*, **58**, 2912-2916.
- Twomey, S., 1974: Pollution and the planetary albedo. *Atmos. Environ.*, **8**, 1251-1256.
- Wang, H., and G. Feingold, 2009: Modeling mesoscale cellular structures and drizzle in marine stratocumulus. Part I: Impact of drizzle on the formation and evolution of open cells. *J. Atmos. Sci.*, **66**, 3237-3256.
- Wood, R., 2009: Private communication.

- Wood, R., T. L. Kubar, and D. L. Hartmann, 2009: Understanding the importance of microphysics and macrophysics for warm rain in marine low clouds. Part II: Heuristic models of rain formation. *J. Atmos. Sci.*, **66**, 2973-2990.
- Wood, R., and couthors, 2008: Open cellular structure in marine stratocumulus sheets. *J. Geophys. Res.*, **113**, D12207, doi:10.1029/2007JD009371.
- Xue, H., G. Feingold, and B. Stevens, 2008: Aerosol effects on clouds, precipitation, and the organization of shallow cumulus convection. *J. Atmos. Sci.*, **65**, 392-406.
- Yum, S. S., J. G. Hudson, 2005: Adiabatic predictions and observations of cloud droplet spectral broadness. *Atmos. Res.*, **73**, 203-223.

Figure Caption

Figure 1. Variation of the liquid water content (L, left), LWP (middle), and cloud thickness (right) with the distance off the California coast into the ocean. The middle and right panels are adapted from Fig. 5b and 5e of Lin et al, respectively for purpose of comparison; the right plot is derived from the data of LWP and cloud thickness according to the method detailed in this comment.

Figure 2. Change of the relative differences in the microphysical properties with the distance off the California coast into the ocean. Note (1) the minus relative difference is shown for the effective radius r_e ; also noted is the different scale used for the drizzle rate.

Table 1 Summary of Quantitative Summer-Winter Differences in Main Properties

Season Property	Summertime	Wintertime	Relative Diff (%)
Cloud fraction	Larger, 74.77%	Smaller, 57.34%	23
Liquid water path	Higher, 70.12	Lower, 40.06	43
Cloud thickness	Similar, 280 m	Similar, 270 m	4
Cloud-base height	Lower, 430 m	Higher, 950 m	-121
Cloud-top height	Lower, 710 m	Higher, 1220 m	-72
LTS*	Stronger, 22 ⁰C	Weaker, 17 ⁰C	23
Inversion strength	Stronger, 7.4 ⁰C	Weaker, 6.0 ⁰C	19
LCL*	Lower, 410 m	Higher, 470 m	-15
Surface-latent heat flux	Smaller, 71 Wm⁻²	Larger, 87 Wm⁻²	-23
SST*	Higher, 19 ⁰C	Lower, 14 ⁰C	26
Liquid water Content	Larger, 0.26 g m⁻³	Smaller, 0.15 g m⁻³	42
Droplet Concentration	Larger, 53 cm⁻³	Smaller, 15 cm⁻³	72
Effective radius	Smaller, 11.4 μm	Larger, 14.6 μm	-28
Drizzle rate	Smaller, 0.67	Larger, 2.38	-255




Figure1

Figure1. Variation of the liquid water content (L, left), LWP (middle), and cloud thickness (right) with the distance off the California coast into the ocean. The middle and right panels are adapted from Fig. 5b and 5e of Lin et al, respectively for purpose of comparison; the right plot is derived from the data of LWP and cloud thickness according to the method detailed in this comment.




Figure 2. Change of the relative differences in the microphysical properties with the distance off the California coast into the ocean. Note (1) the minus relative difference is shown for the effective radius r_e ; also noted is the different scale used for the drizzle rate.